



## Nipox

Liquid Organic Fertilizer Containing  
Vegetable Origin Amino Acid

### PROPERTIES

- NIPOX is a high quality product manufactured using special processes.
- Since it is a product containing high organic carbon, it can be given during the plant development period.
- Increases beneficial microbial activity in the soil due to its high carbon content.
- When mixed with powder fertilizers, it facilitates the uptake of plant nutrients by the roots.
- It does not create salinity in the soil and reduces the salinity effect of other fertilizers due to the organic colloids it contains.
- It is a product containing high levels of amino acids as a result of both the production process and the additive.
- It activates the formation and speed of biochemical events in the plant due to the amino acids it contains.
- It provides an increase in the amount and quality of the product in the plant.
- Increases the durability and shelf life of the product.

Guaranteed Content	(w/w)
Organic Matter	% 52
Organic Carbon	% 19
Total Nitrogen (N)	% 2
Water-soluble Potassium Oxide (K <sub>2</sub> O)	% 4
Free Amino Acids	% 2,5
pH ranger	3,5-5,5





# Nipox

Liquid Organic Fertilizer Containing Vegetable Origin Amino Acid

## APPLICATION DOSES AND PERIODS

PRODUCTS	APPLICATION PERIODS	APPLICATION DOSES	
		LEAF	SOIL
TOMATO, PEPPER	Whenever needed	50-100 CC/ 100 lt water	3-4 lt / da
CUCUMBER, EGGPLANT	Whenever needed	50-100 CC/ 100 lt water	2-3 lt / da
PUMPKIN, MELON, WATERMELON	Whenever needed	50-100 CC/ 100 lt water	3-4 lt / da
BANANA	Whenever needed	100-150 CC/ 100 lt water	4-5 lt / da
CITRUS	Whenever needed	100-150 CC/ 100 lt water	4-5 lt / da
OLIVE	Whenever needed	100-150 CC/ 100 lt water	3-4 lt / da
APPLE, PEAR etc.	After flowering	150-200 CC/ 100 lt water	3-4 lt / da
PEACH, APRICOT, CHERRY, SOUR CHERRY PLUM and POMEGRANATE	After flowering	150-200 CC/ 100 lt water	3-4 lt / da
VINEYARD	After the first leaf is formed	150-200 CC/ 100 lt water	3-4 lt / da
PISTACHIO, HAZELNUT	After flowering	100-150 CC/ 100 lt water	3-4 lt / da
KIWI, FIG	Whenever needed	100-150 CC/ 100 lt water	3-4 lt / da
FRUIT SEEDINGS	Whenever needed	150-200 CC/ 100 lt water	2-3 lt / da
COTTON, SUNFLOWER, CORN etc.	First emergence of the seedlings	100-150 CC/ 100 lt water	3-4 lt / da
TEA PLANT	Whenever needed	100-150 CC/ 100 lt water	3-4 lt / da
PADDY	Whenever needed	100-150 CC/ 100 lt water	2-3 lt / da
LENTIL	First emergence of the seedlings	70-100 CC/ 100 lt water	2-3 lt / da
PEANUT	First emergence of the seedlings	100-150 CC/ 100 lt water	2-3 lt / da
SUGAR BEET	First emergence of the seedlings	100-150 CC/ 100 lt water	3-4 lt / da
CEREALS	In the tillering period of plants	200-250 CC/ 100 lt water	2-3 lt / da
CUT FLOWER	Whenever needed	50-100 CC/ 100 lt water	3-4 lt / da
GREENFIELD PLANTS	Apply 2-3 times as needed	200-300 CC/ 100 lt water	2-3 lt / da

