

Guaranteed Content (w/w)

Organic Matter	% 52
Organic Carbon	% 19
Total Nitrogen (N)	% 2
Water-soluble Potassium Oxide (K 20)	% 4
Free Amino Acids	% 2,5
pH range	3,5-5,5



Nipox

Liquid Organic Fertilizer Containing Vegetable Origin Amino Acid

PROPERTIES

- NIPOX is a high quality product manufactured using special processes.
- Since it is a product containing high organic carbon, it can be given during the plant development period.
- Increases beneficial microbial activity in the soil due to its high carbon content.
- When mixed with powder fertilizers, it facilitates the uptake of plant nutrients by the roots.
- It does not create salinity in the soil and reduces the salinity effect of other fertilizers due to the organic colloids it contains.
- It is a product containing high levels of amino acids as a result of both the production process and the additive.
- It activates the formation and speed of biochemical events in the plant due to the amino acids it contains.
- It provides an increase in the amount and quality of the product in the plant.
- Increases the durability and shelf life of the product

ANTBIO TARIM A.Ş.



Nipox

Liquid Organic Fertilizer Containing Vegetable Origin Amino Acid

APPLICATION DOSES AND PERIODS OF USE			
PRODUCT	PERIODS OF USE	APPLICATION DOSES	
		FOR LEAF	FOR SOIL
TOMATO, PEPPER	Whenever needed	50-100 cc / 100 lt water	3-4 lt / da
CUCUMBER, EGGPLANT	Whenever needed	50-100 cc / 100 lt water	2-3 lt / da
PUMPKIN, MELON, WATERMELON	Whenever needed	50-100 cc / 100 lt water	3-4 lt / da
BANANA	Whenever needed	100-150 cc / 100 lt water	4-5 lt / da
CITRUS	Whenever needed	150-200 cc / 100 lt water	4-5 lt / da
OLIVE	Whenever needed	100-150 cc / 100 lt water	
APPLE, PEAR, etc.	After flowering	150-200 cc / 100 lt water	3-4 lt / da
PEACH, APRICOT, CHERRY, CHERRY, PLUM AND POMEGRANATE	After flowering	150-200 cc / 100 lt water	3-4 lt / da
VINEYARD	After the first leaf is formed	150-200 cc / 100 lt water	
PISTACHIO, HAZELNUT	After flowering	100-150 cc / 100 lt water	3-4 lt / da
KIWI, FIG	Whenever needed	100-150 cc / 100 lt water	3-4 lt / da
FRUIT SEEDINGS	Whenever needed	150-200 cc / 100 lt water	2-3 lt / da
COTTON, SUNFLOWER, CORN, etc.	From the first release	100-150 cc / 100 lt water	3-4 lt / da
TEA	Whenever needed	100-150 cc / 100 lt water	3-4 lt / da
RICE	Whenever needed	100-150 cc / 100 lt water	2-3 lt / da
LENTIL	From the first release	70-100 cc / 100 lt water	2-3 lt / da
PEANUT	From the first release	100-150 cc / 100 lt water	2-3 lt / da
SUGAR BEET	From the first release	100-150 cc / 100 lt water	3-4 lt / da
CEREAL	In the period of reproduction	200-250 cc / 100 lt water	2-3 lt / da
CUT FLORIST	Whenever needed	50-100 cc / 100 lt water	3-4 lt / da
GREENFIELD PLANTS	2-3 times applied whenever needed	200-300 cc / 100 lt water	2-3 lt / da



ANTBİO TARIM A.Ş.